



Flap Necked Chameleon (*Chamaeleo dilepis*)

Widespread and Unique Chameleons

The flap necked chameleon is a medium sized to large chameleon species indigenous to much of Sub-Saharan Africa, and which can range in color from light to dark green, sometimes with lighter or darker dorsal lateral banding. There is also often a pale stripe on their lower flanks, and variable amounts of pale colored stripes, dashes, or spots higher on the flanks. The casque in this species also tends to be low and inconspicuous. This species of chameleon is readily available in captivity, and can do reasonably well provided the correct chameleon care, knowledge, and setup is provided beforehand.

Chameleons in general are a highly diverse and specialized family of Old World lizards with several extremely unique features and adaptations for survival. Perhaps best known are their color and/or pattern changing abilities. The skin and scales of chameleons possess a thin, superficial layer of color changing pigments, with another deeper layer consisting of guanine cells. In order to change their coloration, chameleons rely on changing the space and distance between these guanine cells, thereby changing the wavelengths of light reflected from the skin and scales. While it is popularly believed that chameleons change their colors and patterns for camouflage or crypsis, and while it is true that this has functions in concealing the animal from predators and prey, this is not the primary reason for their color changes. Instead, they will change color as a means of social signaling, to display territoriality and dominance, displaying defensive behaviors, and/or to aid in thermoregulation. Most chameleons also have highly specialized zygodactylus feet and prehensile tails for climbing and gripping branches as well. They also have highly modified and rapidly extrudable sticky tongues used and designed for capturing prey. Chameleons also possess a pair of highly mobile, independently operating eyes, giving them excellent depth perception and motor coordination as well. These are only some of the many unique features chameleons possess, making them popular and unusual pet reptiles to maintain in herpetoculture.

Taxonomy

Life: All living, physical, and animate entities

Domain: Eukaryota

Kingdom: Animalia

Phylum/Sub Phylum: Chordata/Vertebrata

Class: Reptilia

Order: Squamata

Suborder: Lacertilia

<http://www.madisonherps.org>

Infraorder: Iguania

Family: Chamaeleonidae

Genus: Chamaeleo

Species: *Chamaeleo dilepis**

**Taxonomy subject to change and revision.*

Lifespan and Longevity

If provided the proper care, flapneck chameleons can attain longevity of 4 to 8 years on average in captivity.

Distribution and Habitat

The flap necked chameleon is indigenous over a wide range over Sub-Saharan Africa, from Ethiopia and Somalia, south to South Africa in a variety of environments, including coastal forests, moist to dry grasslands and savannahs, woodlands, and brush/scrublands.

Conservation Status

IUCN Red List Least Concern (LC).

Experience Level Required

Novice/Beginner to Intermediate/Moderate.

Size

Adult flap necked chameleons range in size from 9 to 12 inches on average. Some males will reach up to 14 inches or so.

Housing and Enclosure

Housing must be sealed, well ventilated, and escape proof. Hatchling and juvenile chameleons up to around six to eight months can be maintained in roughly a 20 gallon tall enclosure, but be sure to not use too large of enclosures as they can become stressed and have greater difficulty finding their food. Most adult chameleons should be kept in at least a 30 to 50 gallon tall or equivalent sized enclosure. Glass or screen enclosures can be used depending on your household's ambient temperature and humidity. Providing a substrate is optional as chameleons are arboreal and will seldom use it, but substrates that can be provided can include cypress mulch, sphagnum moss, chemical and pesticide free potting soils, or coconut fiber. Chameleons are highly arboreal and specialized lizards that require additional climbing, basking, and hiding opportunities by providing additional vertical artificial foliage, driftwood, branches, logs, and other furnishings. Most chameleons also will not recognize standing water, and should be misted, by hand or with a commercially available misting and/or fogging system to ensure they receive adequate moisture and hydration.

Temperature, Lighting, and Humidity

Provide a lighting and temperature gradient for flap necked chameleons from 75 degrees F to 90 degrees F. A basking temperature of about 95 degrees F. should be provided. Nighttime temperatures can be allowed to drop to 65 to 70 degrees F, and a 12 to 14 hour day/night cycle provided. Use an infrared night time bulb to create nighttime temperature gradients of 75 to 85 degrees F. Providing the correct amounts of UVA/UVB overhead incandescent and florescent lighting, and calcium-to-phosphorus ratios is essential for ensuring the health and overall well-being of flap necked chameleons in captivity. Without UVA/UVB, or adequate amounts of it, they can be susceptible to the abnormal bone growth and development known as Metabolic Bone Disease (MBD), and other health and development maladies. Also be sure to spot clean the enclosure for urates, feces, or uneaten food at least twice per week. Be sure to periodically

<http://www.madisonherps.org>

replace the substrate, clean, and disinfect the enclosure and its furnishings at minimum every 2 to 3 months. More specific lighting, heating, and humidity product suggestions and recommendations that can best suit one's needs, as well as those of one's animals can be given as well.

Feeding, Diet, and Nutrition

Insectivorous; In the wild, flap necked chameleons are primarily insectivorous, feeding on insects and other invertebrates they can locate and capture using their specialized eye-sight and long sticky tongues. In captivity, feed flap necked chameleons a variety of appropriately sized feeder insects such as crickets, roaches, mealworms, superworms, and waxworms. Feeder insects should be gut-loaded in order to increase their optimal nutritional value. Flap necked chameleons also require additional calcium and vitamin D3 supplementation 1 to 2 times weekly or as otherwise directed for optimal health and development. This is in order to prevent Metabolic Bone Disease (MBD) and other growth and nutritional deficiencies. Food can be given by hand or in a small dish or pan mounted off of the enclosure's floor for arboreal animals. Their feeding frequency will depend on the age, size, and overall health of your animal. Use care as to not overfeed them, as obesity and other health related issues can become an issue. More specific dietary and supplementary product suggestions and recommendations that can best suit one's needs, as well as those of one's animals can be given as well.

Handling

Flap necked chameleons are a hardy and easily kept chameleon species. As with most chameleons, however, flap necked chameleons will generally not tolerate frequent handling, and it should be kept to low levels. Should a chameleon be handled, it is important to remember to not forcefully pick up the animal, which may lead to further stress and/or injury. Allow a chameleon to walk onto your hands or gently coax them to do so if possible. Generally, chameleons are better suited as display animals.

****Also be sure to practice basic cleanliness and hygiene associated with proper husbandry after touching or handling any animals or animal enclosures to prevent the possibility of contracting salmonellosis or any other zoonotic pathogens****

Contact

Authored by Eric Roscoe. For any additional questions, comments, and/or concerns regarding this animal, group of animals, or this care sheet, please email and contact the Madison Area Herpetological Society at info@madisonherps.org

Disclaimer: Note that the information provided in these, or any care sheets, are not intended to be all-exhaustive, and further research and care should always be sought and provided when it comes to any species one may prospectively be interested in. These care sheets are also not intended to serve as substitutes for professional veterinary medical care and husbandry should any animal require it. Always seek proper and professional veterinary care for any animal should the need arise, and be prepared ahead of time for any and all husbandry costs and expenses that may occur with any animal beyond the initial purchase. Any animal owned is ultimately a matter of personal/individual care and responsibility.

*Copyright Madison Area Herpetological Society, 2017